

BAG2 Polyclonal Antibody

catalog number: **E-AB-53382**

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

Reactivity	Human
Immunogen	Synthetic peptide of human BAG2
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:20-1:100


Data



Western blot analysis of A549 A431 Hela Jurkat and HEPG2 cell using BAG2 Polyclonal Antibody at dilution of 1:300 Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using BAG2 Polyclonal Antibody at dilution of 1:25(×200)

Observed-MW: Refer to figures

Calculated-MW: 24 kDa



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using BAG2 Polyclonal Antibody at dilution of

1:25(×200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7