

## STK32A Polyclonal Antibody

catalog number: E-AB-19136

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|              |  |
|--------------|--|
| Reactivity   | Human;Mouse  |
| Immunogen    | Fusion protein of human STK32A   |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Antigen affinity purification  |
| Conjugation  | Unconjugated   |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

### Applications

### Recommended Dilution

|     |              |
|-----|--------------|
| WB  | 1:500-1:2000 |
| IHC | 1:100-1:200  |


### Data



Western blot analysis of Mouse heart tissue lysate using STK32A Polyclonal Antibody at dilution of 1:1000

**Observed-MV:Refer to figures**

**Calculated-MV:46 kDa**



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using STK32A Polyclonal Antibody at dilution of 1:95 (x200)

### Preparation & Storage

|          |  |
|----------|--|
| Storage  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

### Background

STK32A (serine/threonine kinase 32A), also known as YANK1, is a 396 amino acid protein that belongs to the superfamily of serine/threonine protein kinases and exists as three isoforms. The gene encoding STK32A maps to human chromosome 5, which is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5 associated and is caused by insertions or deletions within the TCOF1 gene.

### For Research Use Only