



A Reliable Research Partner in Life Science and Medicine

# **ZFP42 Polyclonal Antibody**

catalog number: E-AB-16185

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| $\mathbf{T}$ |      |                           | 4.0          |    |
|--------------|------|---------------------------|--------------|----|
| - 1          | O.C. | cri                       | $\mathbf{n}$ | Λn |
| ய            | CO   | $\mathbf{c}_{\mathbf{I}}$ | JUL          | UЦ |

**Reactivity** Human

**Immunogen** Synthetic peptide of human ZFP42

Host Rabbit Isotype IgG

Purification Affinity purification
Conjugation Unconjugated

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

**WB** 1:200-1:1000

#### Data



Western Blot analysis of A549, K562 and PC3 cell using ZFP42 Polyclonal Antibody at dilution of 1:500

Calculated-MW:35 kDa

### Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

Rex-1 (for reduced expression), also designated zinc finger protein 42 (ZFP42), is an acidic zinc finger protein. Rex-1 contains four repeats of the zinc finger nucleic acid-binding motif and a potential acidic activator domain, suggesting that it is a regulatory protein. Rex-1 localizes to the nucleus and is highly expressed in embryonic stem (ES) and undifferentiated murine F9 teratocarcinoma cells. At the transcriptional level, expression of Rex-1 is reduced when F9 cells are induced to differentiate by the addition of retinoic acid (RA), and Rex-1 repression is enhanced by E1A. The Oct-3/4 transcription factor can either activate or repress the Rex-1 promoter, depending on the cellular environment, while Oct-6 can lower the expression of Rex-1.

### For Research Use Only