Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Monkey ANG1 Antibody Pair Set

Catalog No. E-KAB-0666 Applications ELISA

Synonyms ANG-1;ANGPT1;AGP1;AGPT

Kit components & Storage

Title	Specifications	Storage
Monkey ANG1 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Monkey ANG1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0666)		
		Monkey ANG1 Capture Antibody	Monkey ANG1 Detection Antibody (Biotin)	
Immunogen	Immunogen	Recombinant Monkey ANG1 protein	Recombinant Monkey ANG1 protein	
Information	Swissprot	A0A2K5VW43		
Product details	Reactivity	Monkey	Monkey	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Monkey ANG1 in ELISAs.		

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn

Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Applications

Monkey ANG1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Monkey ANG1 Capture	
Capture		Antibody	10
ELISA Detection	1:1000-1:10000	Monkey ANG1 Detection Antibody (Biotin)	Optical Density
			0.01 100 1000 10000 Monkey ANGI Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Angiopoietins are proteins with important roles in vascular development and angiogenesis. All angiopoietins bind with similar affinity to an endothelial cell-specific tyrosine-protein kinase receptor. The protein encoded by this gene is a secreted glycoprotein that activates the receptor by inducing its tyrosine phosphorylation. It plays a critical role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme and inhibits endothelial permeability. The protein also contributes to blood vessel maturation and stability, and may be involved in early development of the heart. Alternative splicing results in multiple transcript variants encoding distinct isoforms.