

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 700 Anti-Rat CD3 Antibody [G4.18]

Catalog Number: E-AB-F1228UM1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat
Host Mouse

Isotype Mouse IgG3, κ

Clone No. G4.18

Isotype Control Elab Fluor® 700 Mouse IgG3, κ Isotype Control[A112-3] [Product E-AB-F09753M1]

Conjugation Elab Fluor® 700

Conjugation Information Elab Fluor[®] 700 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].

Data



Staining of Rat splenocytes with Elab Fluor[®] 700 Anti-Rat CD3 Antibody[G4.18] (filled gray histogram) or Elab Fluor[®] 700 Mouse IgG3, κ Isotype Control (empty black histogram).

Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8 $^{\circ}\text{C}$ for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names T-cell surface glycoprotein CD3 δ;γ;ε;and ζ chains;CD3 Complex;T3;CD3

Web: www.elabscience.cn

 Uniprot ID
 P19377;Q64159;D4A5M2;

 Gene ID
 25710,300678,315609,25300

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD3 is a complex composed of δ , γ , ϵ , and ζ chains. They are 20-25 kD members of the immunoglobulin superfamily and associated with the T cell receptor (TCR). CD3 is expressed on thymocytes, peripheral T cells, some NK-T cells, and dendritic epidermal T cells. CD3 is involved in antigen recognition, signal transduction, and T cell activation