

Recombinant Mouse IL-11 Protein(Sumo Tag)

Catalog Number: GPEM0336

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	E.coli-derived Mouse IL-11 protein Pro22-Leu199 with an N-terminal Sumo
Calculated MW	32.4 kDa
Observed MW	36 kDa
Accession	P47873
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse IL-11 proteins, 2 µg/lane of Recombinant Mouse IL-11 proteins, was resolved with SDS-PAGE under reducing conditions, showing bands at 36 KD

Background

IL11 is a multifunctional cytokine first isolated in 199 from bone marrow-derived stromal cells. It is a key regulator of multiple events in hematopoiesis, most notably the stimulation of megakaryocyte maturation. IL11 is also known under the names adipogenesis inhibitory factor (AGIF) and oprelvekin. IL11 can improve platelet recovery after chemotherapy-induced thrombocytopenia, induce acute-phase proteins, modulate antigen-antibody responses, participate in the regulation of bone cell proliferation and differentiation, and could be used as a therapeutic for osteoporosis. IL11 stimulates the growth of certain lymphocytes and, in the murine model, stimulates an increase in the cortical thickness and strength of long bones. As a signaling molecule, IL11 has a variety of functions associated with its receptor interleukin 11 receptor alpha, such functions include placentation and to some extent of decidualization.

For Research Use Only