

## FITC Anti-Mouse CD4 Antibody[RM4-5]

**Catalog Number:** GFH1353UC

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

|                                |                                                                                                                                                          |
|--------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Reactivity</b>              | Mouse                                                                                                                                                    |
| <b>Host</b>                    | Rat                                                                                                                                                      |
| <b>Isotype</b>                 | Rat IgG2a, $\kappa$                                                                                                                                      |
| <b>Clone No.</b>               | RM4-5                                                                                                                                                    |
| <b>Isotype Control</b>         | FITC Rat IgG2a, $\kappa$ Isotype Control[2A3] [Product E-AB-F09833C]                                                                                     |
| <b>Conjugation</b>             | FITC                                                                                                                                                     |
| <b>Conjugation Information</b> | FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter). |
| <b>Storage Buffer</b>          | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.                                                                                      |


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu\text{g}/10^6$  cells in 100  $\mu\text{L}$  volume].

### Data



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD8a Antibody and FITC Anti-Mouse CD4 Antibody (Left). Splenocytes are stained with APC Anti-Mouse CD8a Antibody and FITC Rat IgG2a,  $\kappa$  Isotype Control (Right).

### Preparation & Storage

|                 |                                                                                                                                                           |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| <b>Shipping</b> | Ice bag                                                                                                                                                   |

### Antigen Information

|                        |                                                                         |
|------------------------|-------------------------------------------------------------------------|
| <b>Alternate Names</b> | L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4 |
| <b>Gene ID</b>         | 12504                                                                   |

### For Research Use Only

Tel: 400-999-2100

Web: [www.elabscience.cn](http://www.elabscience.cn)

Email: [techsupport@elabscience.cn](mailto:techsupport@elabscience.cn)

Rev. V1.3

## Background

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes and a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a co-receptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosine kinase lck.