A Reliable Research Partner in Life Science and Medicine

Recombinant PTBP1 Monoclonal Antibody

catalog number: E-AB-81602

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Hamster

Immunogen A synthetic peptide of human PTBP1

HostRabbitIsotypeIgGCloneR01-1G2

Purification Affinity Purified


Buffer 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.05% stabilizer and 0.05%

protective protein.

Applications Recommended Dilution

WB 1:1000-1:3000 IHC 1:50-1:100

Data





Western blot detection of PTBP1 in Hela,CHO-K1,Ramos using PTBP1 Rabbit mAb(1:1000 diluted)

Immunohistochemistry of PTBP1 in paraffin-embedded Human tonsil using PTBP1 Rabbit mAb at dilution 1:50

Observed-MW:57 kDa Calculated-MW:57 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA-binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has four repeats of quasi-RNA recognition motif (RRM) domains that bind RNAs. This protein binds to the intronic polypyrimidine tracts that requires pre-mRNA splicing and acts via the protein degradation ubiquitin-proteasome pathway. It may also promote the binding of U2 snRNP to pre-mRNAs. This protein is localized in the nucleoplasm and it is also detected in the perinucleolar structure. Alternatively spliced transcript variants encoding different isoforms have been described.

For Research Use Only

Fax: 1-832-243-6017