# Recombinant Mouse IL-13 Protein(His Tag)

Catalog Number: PDEM100328



Note: Centrifuge before opening to ensure complete recovery of vial contents.

|   |    |  |      | ption |    |              |
|---|----|--|------|-------|----|--------------|
|   | 06 |  | TO T | m     | 11 | $\mathbf{n}$ |
| v |    |  |      |       | ш  | w            |

Species Mouse

**Source** E.coli-derived Mouse IL-13 protein Ser26-Phe131, with an C-teminal His

 Mol\_Mass
 11.5 kDa

 Accession
 P20109

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 80% as determined by reducing SDS-PAGE.

**Endotoxin** < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis

## Data



SDS-PAGE analysis of Mouse IL-13 proteins, 2µg/lane of Recombinant Mouse IL-13 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 30-35 kDa

### **Background**

Interleukin 13 (IL-13) is a single-chain glycosylated polypeptide, which belongs to the IL-13/IL-4 family. IL-13 protein is secreted by many cell types, but especially by T helper type 2 (Th2) cells. IL-13 exerts its effects through a multi-subunit receptor comprising the alpha chain of the IL-4 receptor (IL-4R $\alpha$ ) and at least one of two known IL-13-specific binding chains (IL-13 R $\alpha$ 1 and IL-13 R $\alpha$ 2). As a cytokine, IL-13 protein is critical in regulating inflammatory, immune responses, and diseases. Also, it inhibits the production of pro-inflammatory cytokines and chemokines, and thus down-regulates macrophage activity. IL-13 protein and antibody are more importantly implicated as a central mediator of immunoregulatory processes in various cell types.

### For Research Use Only