Elabscience®


Elab Fluor[®] Violet 450 Anti-Human CD18 Antibody[TS1/18.1.2.11]

Catalog Number: E-AB-F1057Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse lgG1, κ
Clone No.	TS1/18.1.2.11
Isotype Control	Elab Fluor [®] Violet 450 Mouse IgG1, к Isotype Control[MOPC-21] [Product E-AB- F09792Q]
Conjugation	Elab Fluor [®] Violet 450
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with Elab

Fluor[®] Violet 450 Anti-Human CD18 Antibody (filled gray histogram) or Elab Fluor[®] Violet 450 Mouse IgG1 Isotype Control (empty black histogram).

Preparation & Storage		
Storage	Keep as concentrated solution.	
	This product can be stored at 2-8°C for 12 months. Please prote exposure to light and do not freeze.	ected from prolonged
Shipping	Ice bag	
Antigen Information		
Alternate Names	CD18;Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit beta; Complement receptor C3 subunit beta;Integrin beta-2;Itgb2	
Uniprot ID	P05107	
For Research Use Only	/	
Toll-free: 1-888-852-8623 Web: <u>w w w .elabscience.com</u>	Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>	Fax: 1-832-243-601
		Rev. V1.6

Elabscience®

Gene ID Background

3689

CD18 is a 90-95 kD type I transmembrane protein also known as integrin β 2 subunit, LFA-1 β subunit, and β 2 integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.