

Human TARC Antibody Pair Set

Catalog No.	E-KAB-0166	Applications	ELISA
Synonyms	CCL17, ABCD-2, SCYA17, A-152E5.3, Small Inducible Cytokine Subfamily A(Cys-Cys) Member 17		

Kit components & Storage

Title	Specifications	Storage
Human TARC Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze / thaw cycles.
Human TARC Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0166)	
		Human TARC Capture Antibody	Human TARC Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human TARC protein	Recombinant Human TARC protein
	Swissprot	Q92583	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human TARC in ELISAs.	

For Research Use Only

Applications

Human TARC Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images										
ELISA Capture	0.5-4μg/mL	Human TARC Capture Antibody	 <table><caption>Approximate data points from the standard curve</caption><thead><tr><th>Human TARC concentration (pg/mL)</th><th>Optical Density</th></tr></thead><tbody><tr><td>10</td><td>0.05</td></tr><tr><td>100</td><td>0.2</td></tr><tr><td>1000</td><td>1.0</td></tr><tr><td>10000</td><td>5.0</td></tr></tbody></table>	Human TARC concentration (pg/mL)	Optical Density	10	0.05	100	0.2	1000	1.0	10000	5.0
Human TARC concentration (pg/mL)	Optical Density												
10	0.05												
100	0.2												
1000	1.0												
10000	5.0												
ELISA Detection	1:1000-1:10000	Human TARC Detection Antibody (Biotin)											

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

For Research Use Only