Purified Anti-Human CD18 Antibody[TS1/18.1.2.11]

catalog number: E-AB-F1057A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
Isotype	Mouse IgGl, κ
Clone	TS1/18.1.2.11
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze
	to completely remove the stabilizer prior to labeling.
Applications	Recommended Dilution
FCM	$2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$
Data	
Human peripheral blood µg Purified Anti-Huma	10^{2} 10^{3} 10^{4} 10^{5} 10^{6} 10^{2} 10^{3} 10^{4} 10^{5} 10^{6} 10^{10}
followed by FITC-cc Seco	onjugated Goat Anti-Mouse IgG ndary Antibody.
followed by FITC-cc Seco Preparation & Storage	ndary Antibody.
followed by FITC-cc Seco	ondary Antibody. Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
followed by FITC-cc Seco Preparation & Storage	ndary Antibody.

CD18 is a 90-95 kD type I transmembrane protein also known as integrin β 2 subunit, LFA-1 β subunit, and β 2 integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.