

# EIF3K Polyclonal Antibody

catalog number: E-AB-52855

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human;Mouse
<b>Immunogen</b>	Fusion protein of human EIF3K
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Antigen affinity purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

## Applications

## Recommended Dilution

<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:50-1:300


## Data



Western blot analysis of 293T HT29 and A172 cell lysates using EIF3K Polyclonal Antibody at dilution of 1:550

**Observed-MV: Refer to figures**

**Calculated-MV: 25 kDa**



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using EIF3K Polyclonal Antibody at dilution of 1:75 (×200)

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

The 700-kD eukaryotic translation initiation factor-3 (eIF3) is the largest eIF and contains at least 12 subunits, including EIF2S12. eIF3 plays an essential role in translation by binding directly to the 40S ribosomal subunit and promoting formation of the 40S preinitiation complex.

## For Research Use Only