

## ZBTB3 Polyclonal Antibody

catalog number: E-AB-19552

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Human;Mouse
Immunogen	Synthetic peptide of human ZBTB3
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications


### Recommended Dilution

WB	1:500-1:2000
IHC	1:35-1:200

### Data




Western blot analysis of A549 A172 HT-29 HEPG2 and Hela cell lysates using ZBTB3 Polyclonal Antibody at dilution of 1:550



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using ZBTB3 Polyclonal Antibody at dilution of 1:55(×200)

**Observed-MV:Refer to figures**

**Calculated-MV:62 kDa**



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using ZBTB3 Polyclonal Antibody at dilution of 1:55(×200)

### Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

### Background

### For Research Use Only

The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (POxvirus and Zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. ZBTB3 (zinc finger and BTB domain containing 3) is a 574 amino acid protein that contains one BTB (POZ) domain and two C2H2-type zinc fingers. Localized to the nucleus, ZBTB3 is thought to play a role in transcriptional regulation events. The gene encoding ZBTB3 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome.