

IFN- λ 2/IL-28A (C-6His), Human, Recombinant

Cat. No. : PCK160

General Information

Synonyms	Interferon lambda-2; IFN-lambda-2; Cytokine Zcyto20; Interleukin-28A; IL-28A; IFNL2; IL28A; ZCYTO20
Species	Human
Expression host	Human Cells
Sequence	Val26-Val200
Accession	Q8IZJ0
Tag	C-6His
Mol mass	20.6 kDa
Expiration date	12 months

Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Background

IL-28A (Interferon- λ 2; IFN- λ 2), IL-28B/IFN- λ 3, and IL-29/IFN- λ 1 are type III interferons which are distantly related to IL-10 family and type I IFN family Cytokines. Mature human IL-28A is an approximately 22-25 kDa Protein that shares 66% amino acid (aa) sequence identity with mouse and rat IL-28A and shows cross-species activity. It shares 96% and 70% aa sequence identity with human IL-28B and IL-29, respectively. IL-28A promotes the Th1 polarization of dendritic cells in the airway and inhibits Th2 and Th17 mediated inflammation. IL-28A additionally exhibits anti-tumor activity, in part by enhancing IL-12 dependent anti-tumor CTL responses in vivo. In contrast, it is up-regulated in invasive bladder cancer where it promotes tumor cell migration.

SDS-PAGE

