

Purified Anti-Mouse CD161/NK1.1 Antibody[PK136]

catalog number: E-AB-F0987A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse NK1.1 protein
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone	PK136
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM $\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg APC conjugated Anti-Mouse NK1.1 Antibody[PK136] (Right) and 0.2 μg APC conjugated Mouse IgG2a, κ Isotype Control (Left), followed by anti-Mouse CD49b PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

NK-1.1 surface antigen, also known as CD161b/CD161c and Ly-55, is encoded by the NKR-P1B/NKR-P1C gene. It is expressed on NK cells and NK-T cells in some mouse strains, including C57BL/6, FVB/N, and NZB, but not AKR, BALB/c, CBA/J, C3H, DBA/1, DBA/2, NOD, SJL, and 129. Expression of NKR-P1C antigen has been correlated with lysis of tumor cells in vitro and rejection of bone marrow allografts in vivo. NK-1.1 has also been shown to play a role in NK cell activation, IFN- γ production, and cytotoxic granule release. NK-1.1 and DX5 are commonly used as mouse NK cell markers.

For Research Use Only