Elabscience®

Mouse MCP-1 Antibody Pair Set

Catalog No.E-KAB-0089ApplicationsELISASynonymsCCL2, GDCF-2, HC11, HSMCR30, MCAF, MCP1, SCYA2, SMC-CF

Kit components & Storage

Title	Specifications	Storage
Mouse MCP-1 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Mouse MCP-1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0089)	
		Mouse MCP-1 Capture Antibody	Mouse MCP-1 Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Mouse MCP-1 protein	Recombinant Mouse MCP-1 protein
Information	Swissprot	P10148	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse MCP-1 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Mouse MCP-1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Mouse MCP-1 Capture Antibody	
Capture			
ELISA Detection	1:1000-1:10000	Mouse MCP-1 Detection Antibody (Biotin)	Optical Density
			0.01 10 100 100 1000 1000 1000000 10000 10000 10000 10000 10000 100

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4.

For Research Use Only