

## Recombinant Human MICA Protein (His Tag)

**Catalog Number:** PKSH033333

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human MICA protein Glu24-Gln308, with an C-terminal His
<b>Calculated MW</b>	33.9 kDa
<b>Observed MW</b>	60 kDa
<b>Accession</b>	AAH16929.1
<b>Bio-activity</b>	Loaded Human NKG2D-Fc on Protein A Biosensor, can bind Human MICA-His with an affinity constant of 24.6 nM as determined in BLI assay.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

MHC Class I Polypeptide-Related Sequence A (MICA) is a transmembrane glycoprotein that functions as a ligand for human NKG2D. Unlike classical MHC class I molecules, MICA does not form a heterodimer with beta-2-microglobulin. MICA shares 85% amino acid identity with a closely related protein, MICB. MICA acts as a stress-induced self-antigen that is recognized by NK cells, NKT cells, and most of the subtypes of T cells. As a Ligand for the KLRK1/NKG2D receptor, MICA binds to KLRK1 leads to cell lysis. MICA functions as an antigen for gamma delta T cells and is frequently expressed in epithelial tumors. MICA antigens are able to elicit the synthesis of alloantibodies in transplant recipients. Studies have shown that anti-MICA antibodies are associated with acute renal allograft rejection and failure. MICA recognition is involved in tumor surveillance, viral infections, and autoimmune diseases.