

Recombinant Human CD3 ε/CD3E Protein (Fc Tag)

Catalog Number: PKSH032212

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|---|
| Species | Human |
| Source | HEK293 Cells-derived Human CD3 epsilon;CD3E protein Asp23-Asp126, with an C-terminal Fc |
| Calculated MW | 38.7 kDa |
| Observed MW | 40-50 kDa |
| Accession | P07766 |
| Bio-activity | Immobilized Human Human CD3E-Fc at 2µg/ml (100 µl/well) can bind Anti-Human/Monkey CD3E mAb. The ED ₅₀ of Anti-Human/Monkey CD3E mAb is 0.51ug/ml. |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

T-Cell Surface Glycoprotein CD3 ε Chain (CD3ε) is a single-pass type I membrane protein. CD3ε contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3ε is a polypeptide encoded by the CD3E gene on chromosome 11 in humans. The T cell receptor-CD3 complex (TCR/CD3 complex) is involved in T-cell development and several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). TCR/CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017