FUT4 Polyclonal Antibody

catalog number: E-AB-17774



Note: Centrifuge before opening to ensure complete recovery of vial contents.

D		
Description		
Reactivity	Human	
Immunogen	Synthetic peptide of human FUT4	
Host	Rabbit	
Isotype	IgG	
Purification	Antigen affinity purificatio	n
Conjugation	Unconjugated	
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution	
IHC	1:30-1:150	
Data		
cancer tissue using FUT4 Pol	araffin-embedded Human liver yclonal Antibody at dilution of (×200)	Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using FUT4 Polyclonal Antibody at dilution of 1:40(×200)
Preparation & Storage		
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.	
Background		
The product of this gene transfers fucose to N-acetyllactosamine polysaccharides to generate fucosylated carbohydrate		
structures. It catalyzes the synthesis of the non-sialylated antigen, Lewis x (CD15).		

FUT4 (Fucosyltransferase 4) is a Protein Coding gene. Diseases associated with FUT4 include Liver Lymphoma and Colon Adenocarcinoma. Among its related pathways are Mannose type O-glycan biosynthesis and Wnt / Hedgehog / Notch. GO annotations related to this gene include fucosyltransferase activity and alpha-(1->3)-fucosyltransferase activity. An important paralog of this gene is FUT5.

For Research Use Only