

A Reliable Research Partner in Life Science and Medicine

Recombinant ILF3 Monoclonal Antibody

catalog number: AN301758L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;Rat;Mouse

Immunogen Recombinant human ILF3 fragment


HostRabbitIsotypeIgG, κ CloneA466


Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications	Recommended Dilution
WB	1:2000-1:10000
IHC	1:100-1:500
IF	1:50
FCM	1:50-1:100

Data







Western Blot with ILF3 Monoclonal Antibody at dilution of Immunohistochemistry of paraffin-embedded Human kidney 1:10000. Lane 1: A431, Lane 2: Neuro-2a, Lane 3: C6, Lane using ILF3 Monoclonal Antibody at dilution of 1:500.

4: Mouse testis, Lane 5: Rat testis

Observed-MW:90,110 kDa






Immunohistochemistry of paraffin-embedded Human liver using ILF3 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Human testis using ILF3 Monoclonal Antibody at dilution of 1:500.



For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com


Elabscience®


Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Immunohistochemistry of paraffin-embedded Mouse colon using ILF3 Monoclonal Antibody at dilution of 1:500.

Immunohistochemistry of paraffin-embedded Mouse liver using ILF3 Monoclonal Antibody at dilution of 1:500.





Immunohistochemistry of paraffin-embedded Mouse testis using ILF3 Monoclonal Antibody at dilution of 1:500.

Immunohistochemistry of paraffin-embedded Rat cerebrum using ILF3 Monoclonal Antibody at dilution of 1:500.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Interleukin enhancer-binding factor 3 (ILF3) is a double-stranded RNA (dsRNA) binding protein that complexes with other proteins, dsRNAs, small noncoding RNAs, and mRNAs to regulate gene expression and stabilize mRNAs. This protein (NF90, ILF3) forms a heterodimer with a 45 kDa transcription factor (NF45, ILF2) required for T-cell expression of interleukin 2. This complex has been shown to affect the redistribution of nuclear mRNA to the cytoplasm. Knockdown of NF45 or NF90 protein retards cell growth, possibly by inhibition of mRNA stabilization. In contrast, an isoform (NF110) of this gene that is predominantly restricted to the nucleus has only minor effects on cell growth when its levels are reduced. Alternative splicing results in multiple transcript variants encoding distinct isoforms. ILF3 exists main two isoforms NFAR-1 (90 kDa) and -2 (110 kDa). Upon viral infection, ILF3 accumulates in the cytoplasm and participates in the innate antiviral response.

For Research Use Only