

AF/LE Purified Anti-Mouse TCR γ/δ Antibody[UC7-13D5]**Catalog Number:** GF11240**Note:** Centrifuge before opening to ensure complete recovery of vial contents.**Description**

Reactivity	Mouse
Immunogen	Recombinant Mouse TCR γ/δ protein
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone	UC7-13D5
Purification	>98%, Protein A/G purified
Conjugation	None (AF/LE)
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications**Recommended Dilution**

FCM	2 $\mu\text{g/mL}$ (0.5×10^6 - 1×10^6 cells)
------------	---

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR γ/δ belongs to the immunoglobulin superfamily, involved in the recognition of certain bacterial and tumor antigens bound to MHC class I. The TCR γ/δ associates with CD3 and is expressed on a T cell subset found in the thymus, the intestinal epithelium, and the peripheral lymphoid tissues and peritoneum. Most γ/δ T cells are CD4⁻/CD8⁻, some are CD8⁺. T cells expressing the TCR γ/δ have been shown to play a role in oral tolerance, tumor-associated tolerance, and autoimmune disease. It has been reported that γ/δ T cells also play a principal role in antigen presentation. Immobilized UC7-13D5 antibody has been reported to activate TCR- γ/δ -bearing T cells in vitro, and to deplete peripheral TCR- γ/δ -bearing T cells in vivo.