

## Recombinant Human Serglycin/SRGN Protein (His Tag)

**Catalog Number:** PKSH030586

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Serglycin/SRGN protein Met 1-Leu158, with an C-terminal His
<b>Calculated MW</b>	16.1 kDa
<b>Accession</b>	CAA34900.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

SRGN is known as a hematopoietic cell granule proteoglycan. Proteoglycans stored in the secretory granules of various hematopoietic cells has a protease-resistant peptide core, and is vital for neutralizing hydrolytic enzymes. SRGN is associated with the macromolecular complex of granzymes and perforin, which may serve as a mediator of granule-mediated apoptosis. It is required for storage of some proteases in both connective tissue and mucosal mast cells and for storage of granzyme B in T-lymphocytes. SRGN also plays a role in localizing neutrophil elastase in azurophil granules of neutrophils.

### For Research Use Only