ZBTB10 Polyclonal Antibody

catalog number: E-AB-19551



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Synthetic peptide of human ZBTB10

Host Rabbit
Isotype IgG

Purification Antigen affinity purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:50-1:300


Data




Western blot analysis of Jurkat cell lysate using ZBTB10

Polyclonal Antibody at dilution of 1:800

Observed-MV:Refer to figures



Immunohistochemistry of paraffin-embedded Human tonsil tissue using ZBTB10 Polyclonal Antibody at dilution of 1:100(×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using ZBTB10 Polyclonal Antibody at dilution of 1:100(×200)

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

ZBTB10 Polyclonal Antibody

catalog number: E-AB-19551



RINZF, also known as ZBTB10 (zinc finger and BTB domain containing protein 10), is a 847 amino acid protein that contains one BTB/POZ domain and two C2H2-type zinc fingers. Localized to the nucleus, RINZF is believed to play a role in transcriptional regulation. Specifically, RINZF is capable of binding to the CACC element of the Gastrin promoter. In this regard, RINZF competes with Sp1 for CACC binding and interferes with Sp1 transactivation, thereby regulating Gastrin gene expression. The rat RINZF protein shares 98% homology with the human RINZF protein, suggesting that RINZF is a conserved protein. Due to alternative splicing events, two RINZF isoforms exist. In addition, RINZF may be phosphorylated by ATR or ATM upon DNA damage.