

## Purified Anti-Human CD54 Antibody[15.2]

catalog number: E-AB-F10430P

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human CD54 protein
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, κ
<b>Clone</b>	15.2
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Applications

### Recommended Dilution

<b>FCM</b>	2 µg/mL (0.5×10 <sup>6</sup> -1×10 <sup>6</sup> cells)
------------	--

### Data



Human peripheral blood lymphocytes were stained with 0.2 µg Purified Anti-Human CD54 Antibody[15.2] (Right) and 0.2 µg Mouse IgG1, κ Isotype Control (Left), followed

by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD3 FITC-conjugated Monoclonal Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

ICAM-1 is a member of the immunoglobulin superfamily whose expression is upregulated on leukocytes, epithelial cells and resting endothelial cells in response to inflammatory signals. ICAM-1 binds the leukocyte integrins LFA-1 and Mac-1.

### For Research Use Only