

Purified Anti-Human CD54 Antibody[15.2]

catalog number: E-AB-F10430P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD54 protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | 15.2 |
| Purification | >98%, Protein A/G purified |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 μ g/mL (1×10^5 - 5×10^5 cells) |
|------------|---|

Data



Human peripheral blood lymphocytes were stained with 0.2 μ g Purified Anti-Human CD54 Antibody[15.2] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD3 FITC-conjugated Monoclonal Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | Ice bag |

Background

ICAM-1 is a member of the immunoglobulin superfamily whose expression is upregulated on leukocytes, epithelial cells and resting endothelial cells in response to inflammatory signals. ICAM-1 binds the leukocyte integrins LFA-1 and Mac-1.

For Research Use Only