

Purified Anti-Mouse CD209b Antibody[22D1], Functional Grade

catalog number: E-AB-F10220

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD209b protein
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone	22D1
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications


Recommended Dilution

FCM	≤ 0.2 µg per million cells in 100 µL volume
Block	Reported in the literature

Data



Mouse peritoneal macrophages were stained with 0.2 µg Purified Anti-Mouse CD209b Antibody[22D1], Functional Grade(Right) and 0.2 µg Armenian Hamster IgG, κ Isotype Control(Left), followed by Elab Fluor® 647-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody.



Mouse peritoneal macrophages were stained with 0.2 µg Purified Anti-Mouse CD209b Antibody[22D1], Functional Grade(Right) and 0.2 µg Armenian Hamster IgG, κ Isotype Control(Left), followed by Elab Fluor® 647-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.6

CD209B, also known as SIGN-R1, is a mouse C-type lectin receptor predominantly expressed on macrophages in the spleen marginal zone and lymph nodes medulla. CD209B is a mouse homolog of human CD209/DC-SIGN and is involved in innate immune response. CD209B mediates the recognition and uptake of pathogen products, such as lipopolysaccharides (LPS), pneumococcal polysaccharides, and dextrans. CD209B has been demonstrated to facilitate the clearance of encapsulated pneumococcus by directly binding to C1q and activating complement through an immunoglobulin independent pathway.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Leen Moens, et al. Infect Immun. 2007 Dec;75(12):5748-52.