

Recombinant Mouse PD-L1/B7-H1/CD274 Protein (Fc Tag)

Catalog Number: PKSM041256

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|------------------------------------------------------------------------------------------|
| Species | Mouse |
| Source | HEK293 Cells-derived Mouse PD-L1/B7-H1/CD274 protein Phe19-Thr238, with an C-terminal Fc |
| Calculated MW | 51.9 kDa |
| Observed MW | 72-90 kDa |
| Accession | Q9EP73 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. |
| | Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Mouse Programmed cell death 1 ligand 1 (Cd274,PD-L1), is a member of the growing B7 family of immune proteins. It is involved in the costimulatory signal essential for T-cell proliferation and IFN γ production in a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production. B7-H1 has been identified as one of two ligands for programmed death 1 (PD1), a member of the CD28 family of immunoreceptors. B7-H1 is constitutively expressed in several organs such as heart, skeletal muscle. B7-H1 expression is upregulated in a small fraction of activated T and B cells and a much larger fraction of activated monocytes. The costimulatory function of B7-H1 is critical for enhancing maturation and differentiation of T-cells in lymphoid organs. B7-H1 expression is also induced in dendritic cells and keratinocytes after IFN γ stimulation. Interaction of B7-H1 with PD1 results in inhibition of TCR-mediated proliferation and cytokine production. The B7-H1:PD1 pathway is involved in the negative regulation of some immune responses and may play an important role in the regulation of peripheral tolerance.