
## Recombinant Human VEGF-B/VEGFB Protein (Fc Tag)

## Catalog Number: PKSH033474

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Human
HEK293 Cells-derived Human VEGF-B/VEGFB protein Pro22-Ala207, with an N-
terminal Fc
45.7 kDa
40&60 kDa
P49765
Not validated for activity
>95 % as determined by reducing SDS-PAGE.
< 1.0 EU per µg of the protein as determined by the LAL method.
Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
This product is provided as lyophilized powder which is shipped with ice packs.
Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
before lyophilization.
Please refer to the specific buffer information in the printed manual.
Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

## Background

VEGFB, as known as VRF, is a member of the VEGF family of growth factors that share structural and functional similarit y. By alternative splicing, two isoforms of mature VEGF-B containing 167 or 186 amino acid (aa) residues exist. VEGF-B is expressed in most tissues, especially in heart, skeletal muscle and pancreas. The two VEGF-B isoforms have identical amino-terminal cysteine-knot VEGF homology domains but the carboxyl end of VEGF-B167 differs from that of VEGF-B186 by the presence of a highly basic cysteine-rich heparin binding domain. VEGF-B167 and a proteolytically processed form of VEGF-B186 also bind neuropilin-1, a type I transmembrane receptor for semaphorins/collapsins, ligands involved in neuron guidance.

For Research Use Only

Tel:400-999-2100