

A Reliable Research Partner in Life Science and Medicine

## Recombinant Human IL-24 protein(His Tag)

Catalog Number: PKSH034109

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_						
1 B 1	es	0	101	100	41	7
					ш	и

**Species** Human

Source E.coli-derived Human IL-24 protein Gln 52-Leu 206, with an C-terminal His

Calculated MW 19.1 kDa Observed MW 17 kDa Accession O13007

**Bio-activity** Measure by its ability to chemoattract BaF3 cells transfected with human IL-20R alpha

and IL-20R beta. The ED<sub>50</sub> for this effect is <120 pg/mL.

**Properties** 

> 95 % as determined by reducing SDS-PAGE. **Purity** 

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

This product is provided as lyophilized powder which is shipped with ice packs. Shipping

Lyophilized from sterile PBS,pH 8.0. **Formulation** 

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Please refer to the printed manual for detailed information. Reconstitution

## Background

Multifunctional cytokine mainly produced by T-cells that plays a regulatory role in immune response, tissue homeostasis, host defense, and oncogenesis. Possesses antiviral functions and induces the type I intereferon response during influenza infection. Signals through two receptor complexes IL20RA/IL20RB or IL20RB/IL22RA1. In turn, stimulates the JAK1-STAT3 and MAPK pathways and promotes the secretion of pro-inflammatory mediators including IL8 and MMP1. Intracellularly, maintains endoplasmic reticulum homeostasis by restricting the eIF2alpha-CHOP pathway-mediated stress signal. In addition, acts as a quality control mechanism for the ubiquitin proteasome system by alerting the cell to proteasome dysfunction through activation of PKR/EIF2AK2.