IL32 Polyclonal Antibody

catalog number: E-AB-19810



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Synthetic peptide of human IL32
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:40-1:200
Data	
Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using IL32 Polyclonal Antibody at dilution of 1:45(×200)	
Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.
Background	
This gene encodes a member of the cytokine family. The protein contains a tyrosine sulfation site, 3 potential N-	
myristoylation sites, multiple putative phosphorylation sites, and an RGD cell-attachment sequence. Expression of this	

myristoylation sites, multiple putative phosphorylation sites, and an RGD cell-attachment sequence. Expression of this protein is increased after the activation of T-cells by mitogens or the activation of NK cells by IL-2. This protein induces the production of TNFalpha from macrophage cells. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

For Research Use Only