

Recombinant Carbonic Anhydrase VIII/Car8 Monoclonal Antibody

catalog number: **AN300558P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse Carbonic Anhydrase VIII/CA8 protein
Host	Rabbit
Isotype	IgG
Clone	6F2
Purification	Protein A
Buffer	0.2 μ m filtered solution in PBS

Applications

Recommended Dilution


WB	1:500-1:2000
IP	1-4 μ L/mg of lysate

Data



Western Blot with CA8 Monoclonal Antibody at dilution of 1:500 dilution. Lane A: 293T Whole Cell Lysate, Lane B: A549 Whole Cell Lysate, Lysates/proteins at 30 μ g per lane.

Observed-MW:36 kDa
Calculated-MW:33 kDa



Immunoprecipitation analysis using 2 μ L anti-Mouse CA8 Monoclonal Antibody and 15 μ L of 50 % Protein G agarose.

Western blot was performed from the immunoprecipitate using CA8 Monoclonal Antibody at a dilution of 1:200. Lane A:0.5 mg 293T Whole Cell Lysate, Lane B:0.5 mg A549

Whole Cell Lysate

Observed-MW:36 kDa
Calculated-MW:33 kDa

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

The carbonic anhydrases (or carbonate dehydratases) are classified as metalloenzyme for its zinc ion prosthetic group and form a family of enzymes that catalyze the rapid interconversion of carbon dioxide and water to bicarbonate and protons, a reversible reaction that takes part in maintaining acid-base balance in blood and other tissues. The carbonic anhydrase (CA) family consists of at least 11 enzymatically active members and a few inactive homologous proteins. Carbonic anhydrase protein (CA) VIII, which is a member of the CA gene family, has been shown to have no catalytic CA activity and its biological function is still unknown. Increased expression of CA-RP VIII was observed in 78% of colorectal carcinomas. It suggested that CA-RP VIII plays a role in the process of invasion in colorectal cancer.

For Research Use Only