

A Reliable Research Partner in Life Science and Medicine

# PE Anti-Human CD194/CCR4 Antibody[L291H4]

Catalog Number: E-AB-F1366D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ Clone No. L291H4

PE Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792D] Isotype Control

Conjugation

**Conjugation Information** PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


#### **Applications** Recommended usage

**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

# Data



Human peripheral blood lymphocytes are stained with APC Anti-Human CD3 Antibody and PE Anti-Human CD194/CCR4 Antibody[L291H4] (Left). Lymphocytes are stained with APC Anti-Human CD3 Antibody and PE Mouse IgG1, κ Isotype Control (Right).

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

**Shipping** Ice bag

## **Antigen Information**

**Alternate Names** CC-CKR-4;CKR4;CMKBR4;ChemR13;HGCN:14099;K5-5;MGC88293

**Uniprot ID** P51679 Gene ID 1233

# For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

### **Background**

CD194, also known as CCR4, is a CC chemokine receptor. It binds CCL17 and CCL22 and is expressed on a subset of T and B cells, basophils, monocytes, and NK cells. Human Th2 cells are characterized by the expression of CCR4 and CCR8, and these receptors are regulated differently during Th2 development. Human peripheral blood Tregs can be divided into two distinct populations based on the expression of CCR4. Freshly isolated Tregs express CCR4 and presumably represent memory-type Tregs, and CCR4- Tregs require CD3-mediated activation to acquire a regulatory activity. Depletion of CCR4+ T cells leads to Th1-type polarization of CD4+ T cells and augmentation of CD8+ T cell responses to tumor antigens. CCR4 and its ligands are important for the recruitment of memory T cells into the skin in various cutaneous immune diseases.

Fax: 1-832-243-6017