

LDHA Polyclonal Antibody

catalog number: D-AB-10210L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human LDHA protein expressed by E.coli
Host	Rabbit
Isotype	Rabbit IgG
Purification	Antigen Affinity Purification
Buffer	PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4

Applications

Recommended Dilution

WB	1:2500-1:5000
IHC	1:200-1:400


Data




Western blot with Anti LDHA Polyclonal antibody at dilution of 1:2500. Lane 1:HepG2 cell lysate,Lane 2:HeLa cell lysate,Lane 3:Mouse heart tissue lysate,Lane 4:Rat heart tissue lysate.

Observed-MW:36 kDa

Calculated-MW:26,30,36,39 kDa



Immunohistochemistry of paraffin-embedded Rat heart using LDHA Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Mouse heart using LDHA Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.3

LDHA, also named as LDH-M and NY-REN-59, is an enzyme which catalyzes the inter-conversion of pyruvate and L-lactate with concomitant inter-conversion of NADH and NAD⁺. LDHA is found in most somatic tissues, though predominantly in muscle tissue and tumours, and belongs to the lactate dehydrogenase family. It has long been known that many human cancers have higher LDHA levels compared to normal tissues. It has also been shown that LDHA plays an important role in the development, invasion and metastasis of malignancies. Mutations in LDHA have been linked to exertional myoglobinuria. LDHA has some isoforms with MW 26-40 kDa.