A Reliable Research Partner in Life Science and Medicine

Recombinant Human CCL8/MCP-2 Protein (Human Cells, His Tag)

Catalog Number: PKSH032202

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}		crip					
	00	0	РΤ	n	ŤΤ		m
v	\mathbf{c}	v.		w	w	w	ш

Species Human

Source HEK293 Cells-derived Human CCL8; MCP-2 protein Gln24-Pro99, with an C-terminal

His

 Calculated MW
 10.0 kDa

 Observed MW
 13 kDa

 Accession
 A A I26243.1

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA,

pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Human Chemokine (C-C Motif) Ligand 8 (CCL8) is produced by human MG63 osteosarcoma cells. CCL8 shares 62% and 58% amino acid sequence identity with MCP-1 and MCP-3; respectively. All three MCP proteins are monocyte chemoattractants. CCL8 is chemotactic for and activates many different immune cells; including mast cells; eosinophils and basophils; which are implicated in allergic response; and monocytes; T cells; and NK cells that are involved in the inflammatory response. CCL8 elicits its effects by binding to several different cell surface receptors including CCR1; CCR2B and CCR5.

For Research Use Only