

Purified Anti-Human TIGIT Antibody[A15153G]

catalog number: AN009730P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human TIGIT protein
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone	A15153G
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 µg/mL (0.5×10 ⁶ -1×10 ⁶ cells)
------------	--

Data



Human peripheral blood lymphocytes were stained with 0.2 µg Purified Anti-Human TIGIT Antibody[A15153G] (Right) and 0.2 µg Mouse IgG2a, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD56 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

T cell immunoreceptor with Ig and ITIM domains (TIGIT), also known as VSTM3 or WUCAM, is a 26 kD, type I transmembrane protein and is a member of the PVR (poliovirus receptor) family of immunoglobulin-like domain containing proteins. TIGIT is expressed on activated T cells, follicular T helper, memory, and regulatory T cells as well as on NK cells. TIGIT is a negative regulator of NK and T cell activation. Expression of TIGIT is associated with decreased functionality of CD8 T cells in chronic viral infection and tumors. TIGIT also promotes the differentiation of tolerogenic phenotype in dendritic cells with an increased secretion of IL-10 and a diminished production of IL-12.

For Research Use Only