

Recombinant TCF4 Monoclonal Antibody

catalog number: AN302010L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Peptide. This information is proprietary to PTMab.
Host	Rabbit
Isotype	Rabbit IgG
Clone	C331631
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

WB	1:1000
-----------	--------

Data



Western Blot with TCF4 Monoclonal Antibody at dilution of 1:1000. Lane 1: A375, Lane 2: K562, Lane 3: SH-SY 5Y, Lane 4: HeLa

Observed-MW:62 kDa

Calculated-MW:71 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Transcription factor 4 (TCF4), also known as SEF2, ITF2, E2-2, and ME2, binds to the immunoglobulin enhancer Mu-E5/KE5-motif. Involved in the initiation of neuronal differentiation. Activates transcription by binding to the E box (5'-CANNTG-3'). Binds to the E-box present in the somatostatin receptor 2 initiator element (SSTR2-INR) to activate transcription. By similarity. Human TCF4 mRNA expression is particularly high in the brain. Usage of numerous 5' exons of the human TCF4 gene potentially yields in TCF4 protein isoforms with 18 different N-termini. In addition, the diversity of isoforms is increased by alternative splicing of several internal exons. Some isoforms contain a bipartite nuclear localization signal and are exclusively nuclear, whereas distribution of other isoforms relies on heterodimerization partners. TCF4 preferentially binds to either 5'-ACANNTGT-3' or 5'-CCANNTGG-3'. TCF4 has several splicing isoforms that are expressed differentially in tissues and during cancer progression. The scope of molecular weight of several isoforms is about 60-90 kDa.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1