

Recombinant Human HEXA Protein (Subunit A, His Tag)

Catalog Number: PKSH030928

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human HEXA protein Met 1-Thr 529, with an C-terminal His
Calculated MW	59.7 kDa
Observed MW	64 kDa
Accession	AAD13932.1
Bio-activity	Measured by its ability to hydrolyze 4-methylumbelliferyl-N-acetyl-β-D-glucosaminide (4-MU-Glc-NAc). The specific activity is > 1, 250 pmoles/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Wilms' tumor 1-associating protein (WTAP) was previously identified as a protein associated with Wilms' tumor-1 (WT-1) protein that is essential for the development of the genitourinary system. WT1 was originally identified as a tumor suppressor for Wilms' tumor, but it is also overexpressed in a variety of cancer cells. The WTAP-WT1 axis in vascular cells suggest that WTAP is a vital and multifaceted regulator of vascular remodeling. WTAP has been suggested to function in alternative splicing, stabilization of mRNA, and cell growth. Knocking down endogenous WTAP increased Smooth muscle cells (SMCs) proliferation, because of increased DNA synthesis and G(1)/S phase transition, together with reduced apoptosis. These effects could be the result of WTAP suppressing the transcriptional activity of WT1 in SMCs. WTAP may thus also play a role in messenger RNA processing in mammalian cells, either dependent on or independent of its interaction with WT1.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017