PE Anti-Human/Monkey/Mouse Integrin β7 Antibody[FIB504]

Catalog Number: AN00571D


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human;Rhesus;Mouse
Host	Rat
Isotype	Rat IgG2a, κ
Clone No.	FIB504
Isotype Control	PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD3 Antibody and PE Anti-Human/Monkey/Mouse Integrin β7 Antibody[FIB504] (left) or PE Rat IgG2a Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	VI 6T-101;VI A024
Uniprot ID	P26010;P26011

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com

Gene ID	3695;16421
Background	Integrin β7 is a 130 kD glycoprotein also known as integin βp. It is a member of the Ig superfamily. In association with integrin α4 or αE chain, β7 forms α4/β7 or αE/β7 heterodimer. α4/β7 (CD49d/β7, LPAM-1) is expressed on the majority of peripheral lymphocytes, on small subsets of thymocytes, and bone marrow progenitors. LPAM-1 binds to several ligands, VCAM-1, MAdCAM-1 and fibronectin, and is involved in lymphocyte adhesion and some hematopoietic progenitor cells migration. αE/β7 (CD103/β7,αIEL/β7) is expressed on intestinal intraepithelial lymphocytes (IEL), dendritic epidermal T cells, T regulatory cells, a subset of CD8+ T cells in lymph nodes and lamina propria. CD103/β7 complex is thought to play a role in lymphocyte retention via interaction with its ligand E-Cadherin.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com