HSP70 Polyclonal Antibody

catalog number: E-AB-40490



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|---|--|
| Reactivity | Human |
| Immunogen | Synthetic peptide of Human Heat shock 70 kDa protein 1A |
| Host | Rabbit |
| Is otype | IgG |
| Purification | Antigen Affinity Purification |
| Conjugation | Unconjugated |
| buffer | PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4 |
| Applications | Recommended Dilution |
| WB | 1:2000-1:10000 |
| Data | |
| 130 kD→ 95 kD→ 70 kD→ 55 kD→ 40 kD→ | 93T A549 A431 HepG2 |

Western Blot analysis of 293T, A549, A431 and HepG2 cells

 $35 \text{ kD} \rightarrow$ $25 \text{ kD} \rightarrow$ $15 \text{ kD} \rightarrow$

using HSP70 Polyclonal Antibody at dilution of 1:10000.

Observed-MV:70 kDa Calculated-MV:64,70 kDa

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjuction with other heat shock proteins, this protein stabilizes existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins.

For Research Use Only