

A Reliable Research Partner in Life Science and Medicine

FITC Anti-Human CD4 Antibody[SK3]

Catalog Number: E-AB-F1352C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Rhesus; Cynomolgus

Host Mouse

Mouse IgG1, κ Isotype

Clone No. SK3

FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C] Isotype Control

Conjugation

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM**

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Human peripheral blood leucocytes are stained with FITC Anti-Human CD4 Antibody (Left). Leucocytes are stained with FITC Mouse IgG1, κ Isotype Control (Right).

Preparation & Storage

Keep as concentrated solution. Storage

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4 **Alternate Names**

Uniprot ID P01730 Gene ID 920

For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

CD4, also known as T4, is a 55 kD single-chain type I transmembrane glycoprotein expressed on most thymocytes, a subset of T cells, and monocytes/macrophages. CD4, a member of the Ig superfamily, recognizes antigens associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to interact with IL-16.

Fax: 1-832-243-6017