

A Reliable Research Partner in Life Science and Medicine

## **Recombinant ELP3 Monoclonal Antibody**

catalog number: AN301573L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human; Rat; Mouse

Immunogen Recombinant human ELP3 fragment

HostRabbitIsotypeIgG, κCloneA272


Purification Protein Apurified


Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

#### Applications Recommended Dilution

WB 1:500-1:1000 IHC 1:50-1:100 IF 1:50

#### Data






Western Blot with ELP3 Monoclonal Antibody at dilution of 1:1000. Lane 1: HeLa, Lane 2: K562, Lane 3: Mouse brain,


Immunohistochemistry of paraffin-embedded Human tonsil using ELP3 Monoclonal Antibody at dilution of 1:100.

Lane 4: Raw264.7

Observed-MW:62 kDa

Calculated-MW:62 kDa





Immunohistochemistry of paraffin-embedded Mouse testis using ELP3 Monoclonal Antibody at dilution of 1:100.

Immunofluorescent analysis of (100% Ice-cold methanol) fixed HeLa cells using anti-ELP3 Monoclonal Antibody at dilution of 1:50.

Rev. V1.0

#### **Preparation & Storage**

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

### **Background**

#### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Catalytic tRNA acetyltransferase subunit of the RNA polymerase II elongator complex, which is a component of the RNA polymerase II (Pol II) holoenzyme and is involved in transcriptional elongation. The elongator complex is required for multiple tRNA modifications, including mcm5U (5-methoxycarbonylmethyl uridine), mcm5s2U (5-methoxycarbonylmethyl-2-thiouridine), and ncm5U (5-carbamoylmethyl uridine). In the elongator complex, acts as a tRNA uridine(34) acetyltransferase by mediating formation of carboxymethyluridine in the wobble base at position 34 in tRNAs. May also act as a protein lysine acetyltransferase by mediating acetylation of target proteins; such activity is however unclear in vivo and recent evidences suggest that ELP3 primarily acts as a tRNA acetyltransferase. Involved in neurogenesis: regulates the migration and branching of projection neurons in the developing cerebral cortex, through a process depending on alpha-tubulin acetylation. Required for acetylation of GJA1 in the developing cerebral cortex.

For Research Use Only