

EGFR/ErbB1/HER1 (C-6His), Human, Recombinant**Cat. No. : GPCK087****产品信息**

物种	Human
表达宿主	Human Cells
序列信息	Leu25-Ser645
检索号	P00533
标签	C-6His
分子量	69.6 kDa
有效期	12 months
生物活性	Loaded Anti-Human EGFR mAb-Fc on Pro A Biosensor, can bind Recombinant Human EGFR-His with an affinity constant of 7.23 nM as determined in BLI assay.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 8% Sucrose, 50 mM NaCl, 0.05% Tween80, pH 8.0.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

The EGFR subfamily of Receptor tyrosine kinases is composed of EGFR, ErbB2, ErbB3 and ErbB4. The EGFR shares 43%-44% aa sequence identity with the ECD of human EGFR subfamily. All these family members are type I transmembrane glyco Proteins with an extracellular Ligand binding domain. The extracellular Ligand binding domain is containing two cysteine-rich domains separated by a spacer region and a cytoplasmic domain containing a membrane-proximal tyrosine kinase domain. Ligand binding could induce EGFR homodimerization and heterodimerization with ErbB2, resulting in cell signaling, heterodimerization tyrosine phosphorylation and kinase activation. It can bind EGF, amphiregulin, TGF-α, betacellulin, epiregulin, HB-EGF, epigen, and so on. Its signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. EGFR can also be recruited to form heterodimers with the Ligand-activated ErbB3 or ErbB4. EGFR is overexpressed in different tumors. Several anti-cancer drugs use EGFR as target.

网站: www.procell.com.cn

电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



普诺赛® | Procell system

普诺赛® | Procell system

普诺赛® | Procell system

普诺赛® | Procell system

