Recombinant Rat TNF-α Protein(Sumo Tag)


Catalog Number: PDER100248



| Description | |
|----------------|------------------------------------------------------------------------------------------|
| Species | Rat |
| Source | E.coli-derived Rat TNF-a protein Leu80-Leu235, with an N-teminal Sumo |
| Mol_Mass | 30.0 kDa |
| Accession | P16599 |
| Bio-activity | Not validated for activity |
| Properties | |
| Purity | >95% as determined by reducing SDS-PAGE. |
| Endotoxin | < 10 EU/mg of the protein as determined by the LAL method |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5% |
| | Mannitol. |
| Reconstitution | It is recommended that sterile water be added to the vial to prepare a stock solution of |
| | 0.5 mg/mL. Concentration is measured by UV-Vis |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Data



SDS-PAGE analysis of Rat TNF-α proteins, 2µg/lane of Recombinant Rat TNF-α proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 35 kDa

Background

Tumor necrosis factor alpha (TNF-alpha), also known as TNF, TNFA or TNFSF2, is the prototypic cytokine of the TNF superfamily, and is a multifunctional molecule involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. Two receptors, TNF-R1 (TNF receptor type 1; CD120a; p55/60) and TNF-R2 (TNF receptor type 2; CD120b; p75/80), bind to TNF-alpha. TNF-alpha protein is produced mainly by macrophages, and large amounts of this cytokine are released in response to lipopolysaccharide, other bacterial products, and Interleukin-1 (IL-1). TNF-alpha is involved in fighting against the tumorigenesis, thus, is regarded as a molecular insight in cancer treatment.

For Research Use Only