## LMNA Polyclonal Antibody

catalog number: E-AB-31899



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						
	es	OF	ч г	71	IA	m
ய	$\mathbf{c}$	UI	ш.	յս	ΙU	ш

Reactivity Human; Mouse; Rat

**Immunogen** Synthesized peptide derived from human Lamin A/C around the non-phosphorylation

site of Ser392.

Host Rabbit Isotype IgG


Purification Affinity purification Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:100-1:300
IF	1:200-1:1000


#### Data




Western Blot analysis of 3T3 cells using LMNA Polyclonal Immunohistochemistry of paraffin-embedded Human uterus Antibody at dilution of 1:2000.

Observed-MV:74 kDa,65kDa

tissue using LMNA Polyclonal Antibody at dilution of 1:200.





Immunofluorescence analysis of Human liver tissue using LMNA Polyclonal Antibody at dilution of 1:200.

## Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

### Background

## For Research Use Only

# **LMNA Polyclonal Antibody**

catalog number: E-AB-31899



The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants.