

# ZNF85 Polyclonal Antibody

catalog number: E-AB-18071

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Synthetic peptide of human ZNF85
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Antigen affinity purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

## Applications

## Recommended Dilution

<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:20-1:100


## Data



Western blot analysis of Raji and PC3 cell lysates using ZNF85 Polyclonal Antibody at dilution of 1:400

**Observed-MV: Refer to figures**

**Calculated-MV: 69 kDa**



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ZNF85 Polyclonal Antibody at dilution of 1:35 (x200)

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF85 (Zinc finger protein 85), also known as zinc finger protein HPF4 or HTF1, is a member of the ZNF91 family and is thought to be involved in transcriptional regulation. ZNF85 is highly expressed in testicular tissue and localizes to the nucleus. ZNF85 contains sixteen C2H2-type zinc fingers and one KRAB domain through which it is thought to be involved in DNA-binding and transcriptional regulation.

## For Research Use Only