Recombinant Human Interleukin-12/IL-12 Protein

Catalog Number: PKSH033284



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip				
	00	CI	РΤ	n	tπ	Λn
JU.	\mathbf{c}	v.		w	ш	UП

Species Human

 Mol_Mass
 22.5&34.7 kDa

 Accession
 P29459&P29460

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

IL-12 is a heterodimeric pleiotropic cytokine made up of a 40 kDa (p40) subunit and a 35 kDa (p35) subunit. Human and mouse IL-12 share 70% and 60% amino acid sequence identity in their p40 and p35 subunits, respectively. IL-12 is involved in the differentiation of naive T cells into Th1 cells. It is known as a T cell-stimulating factor, which can stimulate the growth and function of T cells. It stimulates the production of interferon-gamma (IFN- γ) and tumor necrosis factor-alpha (TNF- α) from T cells and natural killer (NK) cells, and reduces IL-4 mediated suppression of IFN- γ . T cells that produce IL-12 have a coreceptor, CD30, which is associated with IL-12 activity. IL-12 plays an important role in the activities of natural killer cells and T lymphocytes. IL-12 mediates enhancement of the cytotoxic activity of NK cells and CD8+ cytotoxic T lymphocytes.

For Research Use Only