Elabscience®


Elab Fluor[®] Violet 450 Anti-Human CD15/SSEA-1 Antibody[HI98]

Catalog Number: E-AB-F1079Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse IgM, к
Clone No.	HI98
Isotype Control	Elab Fluor [®] Violet 450 Mouse IgM, κ Isotype Control[MM-30] [Product E-AB-F09782Q]
Conjugation	Elab Fluor [®] Violet 450
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a $450/45$ nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood granulocytes are stained with Elab

Fluor[®] Violet 450 Anti-Human CD15 Antibody (filled gray histogram) or Elab Fluor[®] Violet 450 Mouse IgM, κ Isotype Control (empty black histogram).

Preparation & Storag	ge
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	3-FAL;3-FL;LNFP III;Lewis X;LexSSEA-1;X-hapten;SSEA-1
Uniprot ID	P22083
Gene ID	2526

For Research Use Only

Elabscience®

Background

Elabscience Biotechnology Co., Ltd. A Reliable Research Partner in Life Science and Medicine

CD15 is 3-fucosyl-N-acetyllactosamine (3-FAL), also known as Lewis X, 3-FAL, X-hapten, and SSEA-1. CD15 is expressed on granulocytes and monocytes. It has also been shown to be expressed on Langerhans cells and some malignant cells. CD15 has been implicated in adhesion, as well as chemotaxis, phagocytosis, and bactericidal activity.