

IL-22 (C-6His), Human, Recombinant

Cat. No. : PCK184

General Information

Synonyms	Interleukin-22;IL-22;Cytokine Zcyto18;IL-10-related T-cell-Derived-inducible factor;IL-TIF
Species	Human
Expression host	Human Cells
Sequence	Ala34-Ile179
Accession	Q9GZX6
Mol mass	17.7 kDa
Expiration date	12 months


Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Background

Interleukin-22 (IL-22) is a member of a group of the IL-10 family, a class of potent mediators of cellular inflammatory responses. IL-22 is produced by activated DC and T cells. IL-22 and IL-10 Receptor chains play a role in cellular targeting and signal transduction. It can initiate and regulate innate immune responses against bacterial pathogens especially in epithelial cells such as respiratory and gut epithelial cells. IL-22 along with IL-17 likely plays a role in the coordinated response of both adaptive and innate immune systems. IL-22 also promotes hepatocyte survival in the liver and epithelial cells in the lung and gut similar to IL-10. Biological activity of IL-22 is initiated by binding to a cell-surface complex consisting of IL-22R1 and IL-10R2 Receptor chains. IL-22 biological activity is further regulated by interactions with a soluble binding Protein, IL-22BP. IL-22BP and an extracellular region of IL-22R1 share sequence similarity. In some cases, the pro-inflammatory versus tissue-protective functions of IL-22 are regulated by Cytokine IL-17A.

SDS-PAGE



Procell[®]system

Procell[®]system