

## Elab Fluor® 488 Anti-Human CD57 Antibody[HNK-1]

**Catalog Number:** E-AB-F1067L

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                                |   |
|--------------------------------|---|
| <b>Reactivity</b>              | Human   |
| <b>Host</b>                    | Mouse   |
| <b>Isotype</b>                 | Mouse IgM, κ  |
| <b>Clone No.</b>               | HNK-1   |
| <b>Isotype Control</b>         | Elab Fluor® 488 Mouse IgM, κ Isotype Control[MM-30] [Product E-AB-F09782L]  |
| <b>Conjugation</b>             | Elab Fluor® 488   |
| <b>Conjugation Information</b> | Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter). |
| <b>Storage Buffer</b>          | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.   |

### Applications

### Recommended usage

|            |   |
|------------|---|
| <b>FCM</b> | Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |
|------------|---|

### Data



Staining of normal human peripheral blood cells with APC Anti-Human CD8a Antibody[OKT-8] and Elab Fluor® 488 Anti-Human CD57 Antibody[HNK-1] (left) or Elab Fluor® 488 Mouse IgM, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

|                 |   |
|-----------------|---|
| <b>Storage</b>  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| <b>Shipping</b> | Ice bag   |

### Antigen Information

|                        |                  |
|------------------------|------------------|
| <b>Alternate Names</b> | HNK-1;Leu-7;NK-1 |
| <b>Uniprot ID</b>      | Q9P2W7           |

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.6

**Gene ID**

27087

**Background**

CD57, also known as HNK-1, NK-1, and Leu-7 is a 100-115 kD oligosaccharide antigenic determinant expressed on a variety of proteins, lipids, and chondroitin sulfate proteoglycans. CD57 is expressed on a subset of peripheral blood lymphocytes, including NK cells and CD8+ T cells, and is also expressed on neural cells and striated muscle. CD57 is not expressed on red blood cells, granulocytes, monocytes, or platelets. While the function of CD57 is unknown, binding to L-selectin, P-selectin, and a fragment of laminin suggests that CD57 may be involved in cell-matrix interactions. CD57 is increased in some disease states associated with CD4/CD8 imbalances (AIDS, autoimmune disease, viral infections, and allograft transplants).